



Lichen Sclerosus

What Is Lichen Sclerosus?

Lichen sclerosus is a long-term problem of the skin. It mostly affects the genital and anal areas. Sometimes, lichen sclerosus appears on the upper body, breasts, and upper arms.

Who Gets Lichen Sclerosus?

Lichen sclerosus appears in:

- Women (often after menopause)
- Men (uncommon)
- Children (rare).

What Are the Symptoms?

Early in the disease, small white spots appear on the skin. The spots are usually shiny and smooth. Later, the spots grow into bigger patches. The skin on the patches becomes thin and crinkled. Then the skin tears easily, and bright red or purple bruises are common. Sometimes, the skin becomes scarred. If the disease is a mild case, there may be no symptoms.

Other symptoms are:

- Itching (very common)
- Discomfort or pain
- Bleeding
- Blisters.

What Causes Lichen Sclerosus?

Doctors don't know the exact cause of lichen sclerosus. Some doctors think a too active immune system and hormone problems may play a role. It is also thought that people inherit the likelihood of getting the disease. Sometimes, lichen sclerosus appears on skin that has been damaged or scarred from some other previous injury. Lichen sclerosus is not contagious (it can't be caught from another person).

How Is It Diagnosed?

Doctors can look at severe lichen sclerosus and know what it is. But usually, a doctor takes a small piece of the skin patch (biopsy) and looks at it under a microscope. This allows doctors to make sure that it is not a different disease.

How Is It Treated?

Lichen sclerosus of the genital skin should be treated. Even if it isn't painful or itchy, the patches can scar. This can cause problems with urination or sex. There is also a very small chance that skin cancer may develop in the patches. Surgery is normally a good option for men. Circumcision (removing the foreskin on the penis) is the most widely used therapy for men with lichen sclerosus. The disease usually does not come back. Surgery is normally not a good option for women. When the lichen sclerosus patches are removed from the genitals of women and girls, they usually come back.

Treatment also includes using very strong cortisone cream or ointment on the skin. You put these creams on the patches every day for several weeks. This stops the itching. Then you use the cream or ointment two times a week for a long time to keep the disease from coming back. Treatment does not fix the scarring that may have already occurred. You need regular follow-up by a doctor because using these creams and ointments for a long time can cause:

- Thinning and redness of the skin
- Stretch marks where the cream is applied
- Genital yeast infections.

Lichen Sclerosus

- Sometimes, you don't get better when using the cortisone creams. Some things that can keep symptoms from clearing up are:
- Low estrogen levels
- Infection
- Allergy to the medication.

When creams and ointments don't work, your doctor may suggest:

- Retinoids, or vitamin A-like drugs
- Tacrolimus ointment
- Ultraviolet light treatments (not used on skin of the genitals).

Can People With Lichen Sclerosus Have Sex?

Women with severe lichen sclerosus in the genitals may not be able to have sex. The disease can cause scars that narrow the vagina. Also, sex can hurt and cause the patches to bleed. However, treatment with creams or ointments can help. Women with severe scarring in the vagina may need surgery, but only after lichen sclerosus is controlled with medication.

Is Lichen Sclerosus Related to Cancer?

Lichen sclerosus does not cause skin cancer. However, skin that is scarred by lichen sclerosus is more likely to develop skin cancer. If you have the disease, see the doctor every 6 to 12 months.